

MADE IN FRANCE IN OVER WORKSHOP SINCE 1992

This medical device is intended for patient positioning. This is not a restraint system. Before use, it is important to check the status of the device: stitching, state of the buckles, slings, fabric. Any damage will either be repaired by NAUSICAA Medical, or will result in a disposal.

Y & PELVIC SLIM BELT



TO PREVENT FORWARD SLIDING AND MAINTAIN THE CHEST IN UPRIGHT POSITION FOR ALL CHAIRS EXEPT SHELLS



HOME CARE RANGE 1307 X X XXXXXXX





| | CSYP-80 | WAIST 60/80 cm | Size S | 0 |
|-----------------------|-------------|------------------|---------|---|
| | CSYP-95 | WAIST 75/95 cm | Size M | 0 |
| | CSYP-110 | WAIST 90/110 cm | Size L | 0 |
| | CSYP-130 | WAIST 110/130 cm | Size XL | 0 |
| HOSPITAL RANGE 記述这個人的 | | | | |
| | CSYP-HP-80 | WAIST 60/80 CM | Size S | 0 |
| | CSYP-HP-95 | WAIST 75/95 CM | Size M | 0 |
| | CSYP-HP-110 | WAIST 90/110 CM | Size L | 0 |
| | CSYP-HP-130 | WAIST 110/130 CM | Size XL | 0 |
| | | | | |



2 Allée du Piot - ZA Pôle Actif 0660 Gallargues le Montueux RANCE



Lot

IMPORTANT RECOMMANDATIONS : In order to get a maximum efficiency of these sort of product, it is necessary : - to choose the appropriate size for the patient - to get the best possible adjustment to the patient

These products must not be in direct contact with a wounded skin.

Y & PELVIC SLIM BELT

Compositions :

HOME CARE RANGE

Combination of polyurethan foam and jersey/ cotton polyester Jersey 100% polyester fabric 100% cotton bias 100% polyester strap Nylon buckle Thermo compressed foam

HOSPITAL RANGE

Combination of polyurethan foam and on both faces Jersey/PU coating 100% polyester bias 100% polyester strap Nylon buckle Thermo compressed foam

INSTRUCTIONS FOR USE

• Position the pelvic part on the seat of the chair and let the buckle go out towards the back of the chair.

- Let the patient sit on the chair.
- Position the belt around the patient's waist.
- Close the belt using the velcro strap.

• Place the chest belt being careful that the shoulders pads are positioned on the patient's shoulders.

- Pass the strap of the back part of the belt behind the backrest of the chair.
- Before closing it and without adjusting it, check that the back strap has a female buckle looking downwards and 2 female buckles looking upwards.
- Close the 4 rear buckles, adjust

v 04.03/2017

