

NAUSICAA Médical

www.nausicaa-medical.eu

MADE IN FRANCE IN OVER WORKSHOP SINCE 1992



This medical device is intended for patient positioning. This is not a restraint system.

Before use, it is important to check the status of the device: stitching, state of the buckles, slings, fabric. Any damage will either be repaired by NAUSICAA Medical, or will result in a disposal.

YZA JACKET

FOR ALL CHAIRS







A vest, plus a single size pelvic module, allow you to go from a bust fit to an integral fit according to changing needs without size constraints.

YZA2GBHP-PF







Important Recommandations:

In order to get a maximum efficiency of these sort of product, it is necessary:



- to choose the appropriate size for the patient
- to get the best possible adjustment to the patient

These products must not be in direct contact with a wounded skin.

YZA JACKET

HOSPITAL RANGE
Polyester-coated polyester jersey
lined with polyurethane foam
Bias 91% polyamide - 9% elastane
100% polyester webbing
PVC Shoulder Pad
Nylon buckle

INSTRUCTIONS FOR USE DO NOT USE THIS STRAP IF THE SIZE IS NOT SUITABLE FOR THE PATIENT.

The YZA Jacket comes in 2 modular variants:

1-YZA Bust Jacket (YZA-GB)

a - The pelvic part must be returned inside the vest

b - Put on the bust vest, positioning the shoulder pads well on the shoulders of the person, the buckles behind the back of the chair.

c - Position the belt at the waist of the person.

d - Gird the patient and the back of the chair with the lap strap.

e - Close the loop and adjust the length, taking care not to compress the abdomen.

- f Pass a shoulder strap under the abdominal support strap, connect the loops, and adjust its length.
- 2 -The YZA Integral Jacket (YZA-GI) consists of the YZA Bust Jacket to which is added a Pelvic Support Module.
- a Position the pelvic support module on the chair seat, and secure the 25 mm strap to the back of the chair with minimum play.

b - Sit the patient on the chair and on the strap.

c - Put in place the bust vest, positioning the shoulder pads well on the shoulders of the person, the loops behind the back of the chair.

d - Pass the straps behind the back of the chair.

- e Close the nylon loop of the lap strap, then adjust the length of the strap, being careful not to compress the patient's abdomen.
- f Pass the strap of the pelvic support in the passer of the pelvic part. Close the 50 mm loop of the pelvic support by adjusting its length.

v 05.12/2020

